



Singaporean Journal of Scientific Research(SJSR)

An International Journal (AIJ)

Vol.15.No.1 2023,Pp.34-53

ISSN: 1205-2421

available at :www.sjsronline.com

Paper Received : 24-04-2023

Paper Accepted: 20-05-2023

Paper Reviewed by: 1.Prof. Cheng Yu 2. Dr.KumarRaja

Editor : Dr. R. Rameshkumar

Comparative Anti-oxidative and Acetylcholinesterase inhibition potentials of selected tropical plants in the treatment of Alzheimer's and related cognitive complications

Olatunde Abass Oseni^{1*}, Gbenga Sunday Adeleye²

¹Department of Medical Biochemistry, College of Medicine, Ekiti State University, Ado-Ekiti, Nigeria.

²Department of Physiology, College of Medicine, Ekiti State University, Ado-Ekiti, Nigeria.

ABSTRACT

Acetylcholinesterase inhibitors are important therapeutic agents in Alzheimer's and other cognitive complications. The study aimed to evaluate acetylcholinesterase inhibition, antioxidant activity and other biochemical indices of the three solvent-extracts of five plant species from South-western Nigeria.

The leaves were extracted with solvents; phytochemical screening and in-vitro antioxidant properties of the extracts were determined. Rats weighing between 140 to 150g were divided into groups comprising of normal control, 2,2-dichlorovinyl dimethylphosphate (DDVP), aqueous, methanol and ethanol groups for each of the five plants. The rats were induced with 3.3mg/kg body weight of DDVP for two weeks with the aim of inducing cognitive complications while the induced animals were treated with 3.3mg/kg body weight of the extracts of the five plant species for another two weeks. The rats were anaesthetized and quickly dissected individually to collect serum and organs homogenates to evaluate the studied parameters using standard procedures. The results obtained from the study clearly demonstrated that the three extracts of the five plants show

powerful in-vitro and in-vivo antioxidant properties, impressive anti-lipid peroxidation potentials and promising acetylcholinesterase inhibition activity. These plants solvents extracts could be locally utilized in drug developments to prevent and cure the consequences of Alzheimer disease (AD) and associated disorders.

Keywords:Alzheimer disease, cholinergic functions, antioxidant property, acetylcholinesterase inhibitor, drug development

1. INTRODUCTION

Alzheimer's disease is characterized by selective neuronal cell death, the presence of extra cellular amyloid deposits in the core of neurotic plaques and the formation of intra neuronal neurofibrillary tangles in the brain of afflicted individuals (Dhanasekaran *et al.*, 2015). According to the cholinergic hypothesis, the inhibition of acetylcholinesterase (AChE), an enzyme that catalyzes acetylcholine hydrolysis, increases the levels of acetylcholine in the brain, thus improving cholinergic functions in AD patients (Murray *et al.*, 2013). The cholinergic hypothesis states that decreased cholinergic transmission plays a major role in the expression of cognitive, functional and possibly behavioral symptoms in AD (Hau *et al.* 2008; Volpicelli-Daley *et al.*, 2003). The cholinergic hypothesis rests on pathological, biochemical and pharmacological observations. Oxidative stress is also implicated in the development of many neurodegenerative diseases including Parkinson's disease (PD), Huntington's disease, amyotrophic lateral sclerosis and Alzheimer's disease (AD)⁵. One of the important strategies for treating of AD is to maintain the levels (Helena *et al.*, 2012) of acetylcholine through the inhibition of acetylcholinesterase (AChE) (Lahiri *et al.*, 2002). The acetylcholinesterase enzyme (AChE) is an attractive target for the rational drug design and for the discovery of mechanism-based inhibitors because of its role in the hydrolysis of the neurotransmitter acetylcholine (ACh). AChE inhibitors are the most effective approach to treat the cognitive symptoms of Alzheimer disease (AD) (Kalauni *et al.*, 2002; Atta-ur-Rahman *et al.*, 2004) and other possible therapeutic applications in the treatment of Parkinson's disease, senile dementia, and ataxia, among others (Ahmad *et al.*, 2003). AChE inhibitors as eserine, tacrine, donepezil, rivastigmine, and galanthamine are the only drugs currently approved for the treatment of AD; however, these drugs are known to have limitations for clinical use due to their short-half-lives and/or unfavorable side-effects (Sunget *et al.*, 2002). Dichlorvos is a synthetic organic chemical used as an insecticide. Dichlorvos does not occur naturally in the environment but is manufactured by industry. Dichlorvos is an insecticide used on crops, animals, and in pest-strips. Acute (short-term) and chronic (long-term) exposures of humans to dichlorvos results in the inhibition of an enzyme, acetylcholinesterase, with neurotoxic effects including perspiration, vomiting, diarrhea, drowsiness, fatigue, headache, and at high concentrations, convulsions, and coma (TMI, 1989). Since the synthetic drugs have undesirable side effects or contraindications, the World Health Organization (WHO) has recommended the evaluation of traditional plants treatments for diabetes (Day, 1998). Medicinal plants are widely used worldwide to address a variety of health problems. About 25 to 50% of current pharmaceuticals are derived from plants (Cowan, 1999; Goh *et al.*, 1995). Plants are rich in a wide variety of phytochemical metabolites which are divided into two groups: Primary and Secondary metabolites. Primary metabolites consist of common sugars, amino acids, proteins and chlorophylls, while Secondary metabolites include glycosides, alkaloids, saponins, phenolic

compounds, terpenes steroids and anthraquinones (Habtermariam *et al.*, 1993). These plants have been claimed to be used in the treatments of diseases, but its AChE inhibition activity has not been studied so far. Hence the main objective of the present study is to evaluate AChE inhibition, antioxidant activity and other biochemical indices of the three solvent-extracts of five plant species [Scent leaf, ewe efirin (*Ocimumgratissimum*), Wind braker leaf, ewe igunnu (*Polyalthia longifolia* (Sonn.)), Pawpaw leaf, ewe ibepe (*Carica papaya*), African mahogany leaf, ewe apa (*Afzeliaafricana*) and Goat weed leaf, ewe imi-esu (*Ageratum conyzoides*)] from South-western Nigeria.

2. MATERIALS AND METHODS

Plants samples collection, authentication and preparation

The various plants leave of *Ocimumgratissimum*(UHAE 2020051), *Polyalthia longifolia* (Sonn.) (UHAE 2020054), *Carica papaya* (UHAE 2020048), *Afzelia Africana* (UHAE 2020046) and *Ageratum conyzoides* (UHAE2020047) were sourced from various locations in Ado-Ekiti, Ifaki-Ekiti, Iworoko-Ekiti in Ekiti State and Ibadan in Oyo State Nigeria by detaching large quantities of leaves from their various trees of the plants. The leaves were however authenticated at Department of Plant Science of Ekiti State University, Ado-Ekiti, Nigeria and respective Herbarium numbers obtained.

The leaves were rinsed in water to remove any form of dirt's. The rinsed leaves were spread in a room for some days till they were completely air dried. The air-dried leaves were pulverized with electric blender to increase the surface area and to hasten the process of extraction.

Extraction with water, methanol and ethanol

The individual powdered leaves were divided and measured into three parts using standard weighing balance. 200g each of the powdered leaves were extracted with 500mL each of water, methanol and ethanol inside stoppered glass bottles with continuous shaking overnight. The extracted solutions were evaporated to obtain the respective extracts. The resulting extracts were weighed and the percentage solvents extracts yields recorded.

Experimental Animals

The Wistar albino rats weighing between 140g to 150g were obtained from Animal House of College of Medicine, Ekiti State University, Ado-Ekiti, Nigeria.

Animal Care and Management

Eighty-five (85) Wistar albino rats weighing between 140-150g obtained from Animal House of Ekiti State University College of Medicine, Ado-Ekiti, Nigeria were grouped into five groups of five animals each. They were housed in plastic cages under standard laboratory conditions of natural light/dark cycle at room temperature and humidity. The rats were fed on standard rat pellets which were procured once from (Top feeds, Nigeria) and given drinking water *ad libitum*. All animals were handled in accordance with the Guide for the care and use of laboratory animals as detailed in the International Animal Care and Use Committee (IACUC, 2010)

Experimental design and animal assay

The rats were induced with 3.3mg/kg body weight of 2,2-Dichlorovinyl dimethylphosphate (DDVP) solution for fourteen days except group 1. The induced rats were then treated with 3.3mg/kg body weight solution of aqueous, methanolic and ethanolic extracts of the various plants for the next fourteen days.

Group1 Normal Control (animals given water and rats feed *ad libitum*)

Group2 DDVP induced control (animals were administered orally with 0.5mL of 3.3mg/kg body weight of DDVP solution)

Group3 (3i-3v) DDVP induced animals + 0.5mL of 3.3mg/kg body weight of aqueous solution of each extract of the plants.

Group4 (4i-4v) DDVP induced animals + 0.5mL of 3.3mg/kg body weight of methanolic solution of each extract of the plants.

Group5 (5i-5v) DDVP induced animals + 0.5mL of 3.3mg/kg body weight of ethanolic solution of each extract of the plants.

After the experiment, the rats in all the groups were anaesthetized with chloroform individually and quickly dissected to collect blood into the sample bottles and latter centrifuged at 3000rpm to obtain serum; while the organs (brain, liver and heart) were removed, weighed and placed on ice-bath. 10% of each organ homogenate was then prepared respectively in 6.7mM potassium phosphate buffer, (pH 7.4) using the electrically top driven homogenizer.

The individual organ homogenates were centrifuged at 3,000rpm for 10 minutes at 4°C to obtain clear supernatants which were stored at 8°C and used for measurement of the studied biochemical parameters.

The antioxidants enzymes activities [Catalase (CAT), superoxide dismutase (SOD), Glutathione-S-transferase (GST), Glutathione reductase (GR) and Glutathione peroxidase (GPx)]; reduced glutathione (GSH) were determined by the methods described by (Chance, Maehly, 1955; Von Euler, Josephson, 1972; Misra, Fridovich, 1972; Habiget *al.*, 1974; Carlberg, Mannervik, 1985; Mohandaset *al.*, 1984; Jollowet *al.*, 1974) respectively. The lipid peroxidation was done by measuring the TBARS in accordance with the modified method of Utley et al., (1967). The Sigma-Aldrich, (2013) assay Kit was employed to determine the acetylcholinesterase activity, where one unit of AChE is the amount of enzyme that catalyzes the production of 1.0 µmole of thiocholine per minute at room temperature at pH 7.5 with the aim of evaluating the inhibition of acetylcholinesterase activity. This kit has a linear range of 10–600units/L of AChE activity.

Statistical Analysis

The mean triplicate results were determined and represented in charts as presented in the various figures.

RESULTS

The phytochemical compositional screening can be seen in Figure 1 showing the leaves of the five plants containing important compounds of medicinal significant at varying intensity in all the solvents extracts as studied.




FIGURE 1-Phytochemical compositional evaluation of leaves extracts of five Nigerian medicinal plants. Less than 100 implies traces, 100 = “+”, 200 = “++”, 300 = “+++”

Similarly, Figures 2 represents the (% ABTS; % DPPH; % H₂O₂; % Nitric oxide and % Super oxide) scavenging activities and concentrations of Total flavonoids and Total phenol of leaves

extracts of plants, the scavenging activities of each plant extract increased in dose dependent manner.




FIGURE 2- Some in-vitro antioxidants activities of leaves extracts of five Nigerian medicinal plants

The effects of various extracts of the plants on the activity of Catalase as seen in Figure 3, the serum, liver, heart and brain tissues of induced and treated animals showing all the plants causing restored activity of the enzyme after treatments.




FIGURE 3-Effects of various extracts of some Nigerian medicinal plants on the activity of Catalase (u/ml/min) enzyme in 2,2-dichlorovinyl dimethylphosphate induced Wistar albino rats

The percentage superoxide dismutase activity as presented in Figure 4 showed the effects of various extracts of the plants indicating all the solvents extracts display increased SOD percentage activity when compared with the DDVP induced group.




FIGURE 4-Effects of various extracts of some Nigerian medicinal plants on the activity of Superoxide Dismutase (% SOD activity) in 2,2-dichlorovinyl dimethylphosphate induced Wistar albino rats

The glutathione peroxidase activity of various extracts of plants from Figure 5 showed that the extracts caused a reversal effect to normal control group when compared with the DDVP induced group.




FIGURE 5-Effects of various extracts of some Nigerian medicinal plants on the activity of Glutathione peroxidase (Gpx U/L) in 2,2-dichlorovinyl dimethylphosphate induced Wistar albino rats.

Figure 6 showed the Glutathione transferase activity in DDVP induction and treatment with various extracts of the plants, the extracts caused enhanced activity of the enzyme on the treated groups

G. Transferase Activity




FIGURE 6-Effects of various extracts of some Nigerian medicinal plants on the activity of Glutathione transferase ($\mu\text{mol}/\text{min}/\text{ml}$) in 2,2-dichlorovinyl dimethylphosphate induced Wistar albino rats

The activity of glutathione reductase in Figure 7 revealed the improved activity on treatment with the various extracts of the plants.

G. Reductase Activity




FIGURE 7-Effects of various extracts of some Nigerian medicinal plants on the activity of Glutathione reductase (U/L) in 2,2-dichlorovinyl dimethylphosphate induced Wistar albino rats

The concentration of reduced glutathione from Figure 8 showed that the various extracts of the plants caused increased concentration of GSH in the depleted concentration of the DDVP induction group.




FIGURE 8-Effects of various extracts of some Nigerian medicinal plants on the concentration of Reduced glutathione (GSH mM)in 2,2-dichlorovinyl dimethylphosphate induced Wistar albino rats

From Figure 9, the concentration of MDA was grossly reduced from the DDVP induced group when compared to the solvent's extracts treated groups.




FIGURE 9-Effects of various extracts of some Nigerian medicinal plants on the MDA (nmolTBARS/h/mg tissue protein) in 2,2-dichlorovinyl dimethylphosphate induced Wistar albino rats

The inhibition of the Acetylcholinesterase activity observed in Figure 10 showed that the various extracts of the plants caused substantive activity inhibition from the DDVP induction group.

Acetylcholinesterase Activity




FIGURE 10- Effects of various extracts of some Nigerian medicinal plants on the Inhibition of Acetylcholinesterase activity(U/mL) inthe Serum and Brain of 2,2-dichlorovinyl dimethylphosphate induced Wistar albino rats.

DISCUSSION

The qualitative phytochemical composition of the five plants leaves solvents extracts were evaluated to assess the presence of bioactive components. All the studied plants contained few and/or all the tested phytochemicals (alkaloids, saponins, tannins, phenolic compounds, phlobatannin, steroids, anthraquinone, terpenoids and cardiac glycosides) as presented in Figure 1. All the studied phytochemicals were present in either aqueous, methanolic, ethanolic and in all extracts of the plants except anthraquinone that was present in only aqueous extracts of *Carica papaya*, *Afzelia Africana* and *Ageratum conyzoides*. These secondary plant metabolites have been reported to contain antibacterial potency and have been actively used or in combination with antibiotics in the therapy of bacterial infections (Liu et al., 2001; Deba et al., 2008). The demand in the use of natural products as antioxidants and antimicrobial compounds has led to investigate various extracts from many plants, this has recently been of great interest in both research and the food industry, because of growing

tendency and possibility to replace synthetic antioxidants and antimicrobials compounds with natural ones³⁷. Many other researchers have reported diverse plants and plants parts solvents extracts containing arrays of these phytochemicals worldwide (Vaghasiyaet al., 2011; Oseni, Williams, 2018). The in-vitro antioxidants scavenging potentials of (%H₂O₂; % ABTS; % Nitric oxide, % DPPH;% Super oxide scavenging activities and concentrations of total phenol and total flavonoids were investigated as reported in Figure2. Reactive oxygen species (ROS) is a term that encompasses all highly reactive, oxygen containing molecules, including free radicals like hydroxyl radical, the superoxide anion radical, hydrogen peroxide, singlet oxygen, nitric oxide radical, hypochlorite radical, and various lipid peroxides with capability of reacting with membrane lipids, nucleic acids, proteins and enzymes, and other small molecules, resulting in cellular damage (Krishnamurthy, Wadhvani, 2012). Antioxidant compounds may function as freeradical scavengers, initiator of the complexes of prooxidant metals, reducing agents and quenchers of singlet oxygen formation. The in-vitro antioxidants scavenging potentials results obtained in this study however corroborated the observations of earlier investigations by other researchers worldwide revealing plants as major sources of antioxidants (Al-Mustafa, Al-Thunibat, 2008; Nilima, Hande, 2011; Ashafaet al., 2010; Ilahiet al., 2013).Antioxidant's enzymes (CAT, SOD, GPx, GR, GST) and GSH were also investigated as presented in Figures 3.0-8.0. The results showed the plants extracts exhibited high antioxidant enzyme activities and promising concentration of non-enzyme antioxidant compound (GSH). Antioxidants are molecules that prevent oxidation or inactivates the reactive oxygen species and thus prevent oxidative damage to the cells and body tissues (Aliyuet al., 2019).It has been reported that catalase (CAT) is an enzymatic antioxidant widely distributed in all animal tissues, decomposes hydrogen peroxide and protects the tissue from highly reactive hydroxyl radicals (Nandyet al., 2012). Its activity varies greatly from tissue to tissue, the highest activity is found in liver and kidney, whereas the lowest activity has been reported in the connective tissue (Naazeriet al., 2014). It is evidenced in this study that the five plants and the three extracts enhanced the activity of CAT in serum and all the organs studied, this is in agreement with what was obtained by (Merghemet al., 2019; Doungueet al., 2018).The enhanced catalase activity could be due to the presence of bioactive compounds present in these plants extracts that would give their protons to stabilize radicals formed during stress. Administration of DDVP significantly decrease SOD activity in the studied organs while the five plants solvents extracts caused a concomitant increased activity in the treated groups. Similar trends were also observed in GPx, GR and GST activities. The concentration of reduced glutathione (GSH) was also observed to be grossly reduced by the administration of DDVP but treatments with the plants solvents extracts produced upward effects by restoring the GSH concentrations in the serum and the studied organs. In all the antioxidants investigations, the plants solvent extracts reduced the effects of DDVP and hence reversed the oxidative stress expressed by DDVP. From Figure 9 showing the effects of various extracts of plants on the MDA in DDVP induction, there was increased MDA concentrations in both serum and the studied organs indicating lipid peroxidation which implicates oxidative stress. On treatments, all the plants extract showed protective role by decreasing the lipid peroxidation level in DDVP induced animals and increased activities of enzymatic (CAT, SOD, GPx, GR and GST) antioxidants and the level of non-enzymatic (GSH) antioxidant. The results obtained in this study however corroborates those reported by other researchers in their various works on different plants extracts and effects on oxidative stress (Ojoet al., 2014a; Ojoet al., 2014b; Yadavet al., 2012; Ahlamet et al., 2014; Armelleet al., 2018).Figure 10 showed the

results of effects of various extracts of the five plants on the inhibition of acetylcholinesterase activity in the serum and brain of DDVP induced animals. The results obtained in this study showed that DDVP induction of 3.3mg/kg body weight for 14 days significantly elevated AChE in both serum and brain tissues, it has also been reported by other researchers that DDVP caused increase AChE activity and oxidative stress (Ojoet *al.*, 2014a; Maysaaet *al.*, 2016; Yadavet *al.*, 2012;). It has been reported that AChE interact with amyloid β and promotes the formation of amyloid fibril through a pool of amino acids situated in proximity of the peripheral anionic site (PAS) of the enzyme resulting in neurocognitive impairment (Anandet *al.*, 2012). Several studies have revealed that AChE inhibitors not only facilitate transmission of the cholinergic system, but also interfere with the synthesis, aggregation and deposition of toxic amyloid β (Anandet *al.*, 2012).In addition, dysregulation of AChE levels has also been reported in myasthenia gravis, glaucoma, Lewy bodies and Parkinson's disease (Colovicet *al.*, 2013).The observations in this study however corroborated the earlier similar works ofDoungueet *al.*, (2018)in neuroprotective effect and antioxidant activity of *Passiflora edulis* fruit flavonoid fraction, aqueous extract, and juice in aluminum chloride-induced Alzheimer's disease rats; Mahdy *et al.*, (2012) in effect of some medicinal plant extracts on the oxidative stress status in Alzheimer's disease induced in rats andNwiduet *al.*, (2017)in anti-acetylcholinesterase activity and antioxidant properties of extracts and fractions of *Carpolobia lutea*.

CONCLUSION

The inhibition of the AChE activity by the plants solvents extracts increased the acetylcholine concentration of the serum and the brain tissues with positive impact on the cognitive function, hence this study clearly showed that these plants solvents extracts possessed AChE inhibition and antioxidant properties thus could reverse DDVP-induced cognitive dysfunction like Alzheimer's disease and other related cognitive complications.

ACKNOWLEDGEMENT

Mr. Olayinka Okoh of Department of Chemical Sciences, Faculty of Science and Science Education, Anchor University Lagos is hereby acknowledged for his inputs in results presentations.

REFERENCES

- [1] Ahlamet AMA, Ali AA, Mohammed AAS. Effect of White Tea Extract on Antioxidant Enzyme Activities of Streptozotocin –Induced Diabetic Rats. Egypt. Acad. J. Biolog. Sci. 2014; 6(2): 17-30
- [2] Ahmad W, Ahmad B, Ahmad M, Iqbal Z, Nisar M, Ahmad M. *In vitro* inhibition of acetylcholinesterase, butyrylcholinesterase and lipoxxygenase by crude extract of *Myricaria elegans* Royle. *J Biol Sci.* 2003; 11: 1046-49.
- [3] Aliyu AB, Achika JI, Adewuyi JA, Gangas P, Ibrahim H, Oyewale AO. Antioxidants from Nigerian Medicinal Plants: What is the Evidence? Lipid Peroxidation Research, Mahmoud Ahmed Mansour, IntechOpen, 2019, DOI: 10.5772/intechopen.84454

- [4] Al-Mustafa AH, Al-Thunibat OY. Antioxidant Activity of Some Jordanian Medicinal Plants Used Traditionally for Treatment of Diabetes. *Pakistan Journal of Biological Sciences*. 2008; 11 (3): 351-58.
- [5] Anand P, Singh B, Singh N. A review on coumarins as acetylcholinesterase inhibitors for Alzheimer's disease. *Bioorg. Med. Chem*. 2012; 20: 1175–80
- [6] Armelle DT, Lauve RYT, Nole T, Protus AT, Jules-Roger K, Gabriel AA. *Costusafer* Protects Cardio-, Hepato-, and Reno-Antioxidant Status in Streptozotocin-Intoxicated Wistar Rats. *Biomedical Research International*. Volume 2018 |Article ID 4907648 | 9 pages
- [7] Ashafa AO, Grierson DS, Afolayan AJ. *In Vitro* Antioxidant Activity of Extracts From the Leaves of *Felicia Muricata* Thunb. an Underutilized Medicinal Plant in the Eastern Cape Province, South Africa. *Afr J Tradit Complement Altern Med*. 2010; 7(4): 296–302.
- [8] Atta-ur-Rahman, Wahab AT, Nawas SA, Choudhary MI. New cholinesterase inhibiting bisbenzylisoquinoline alkaloids from *Cocculus pendulus*. *Chem Pharm Bull*. 2004; 52: 802-06.
- [9] Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *Lebensmittel-Wissenschaft und-Technologie*. 1995; 28: 25–30.
- [10] Carlberg I, Mannervik B. Glutathione reductase. *Meth. Enzymol*. 1985; 113: 485- 90.
- [11] Chance B, Maehly AC. Assay of catalase and peroxidase. *Meth. Enzymol*. 1955; 2: 764-75.
- [12] Colovic MB, Krstic DZ, Lazarevic-Pasti TD, Bondzic AM, Vasic VM. Acetylcholinesterase inhibitors: pharmacology and toxicology. *Curr. Neuropharmacol*. 2013; 11: 315–35
- [13] Cowan MM. Plants products as anti-microbial agents. *Clin. Microbiol. Rev*. 1999; 12: 564–82.
- [14] Day C. Traditional plant treatments for diabetes mellitus: pharmaceutical foods. *Br. J. Nutr*. 1998; 80: 5–6
- [15] Deba F, Xuan TD, Yasuda M, Tawatu S. Chemical composition and antioxidant, antibacterial and antifungal activities of the essential oils from *Bidens pilosa* Linn. var. *Radiata*. *Food Control*. 2008; 19: 346-52.
- [16] Dhanasekaran S, Perumal P, Palayan M, In-vitro Screening for acetylcholinesterase enzyme inhibition potential and antioxidant activity of extracts of *Ipomoea aquatica* Forsk: therapeutic lead for Alzheimer's disease. *J. Appl. Pharm. Sci*. 2015; 5(02): 12-16.
- [17] Doungue HT, Kengne APN, Kuate D. Neuroprotective effect and antioxidant activity of *Passiflora edulis* fruit flavonoid fraction, aqueous extract, and juice in aluminum chloride-induced Alzheimer's disease rats. *Nutrire*. 2018; 43: 23.
- [18] Goh SH, Chuah CH, Mok JSL, Soepadmo E. *Malaysian Medicinal Plants for the Treatment of Cardiovascular Diseases*. Malaysia: Pelanduk Publication; 1995. 162.
- [19] Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. Analysis of nitrate, nitrite, and [15N] nitrate in biological fluids. *Anal Biochem*. 1982; 126(1):131-38.
- [20] Habig WH, Pabst MS, Jekpoly WB. Glutathione transferase: A first enzymatic step in mercapturic acid formation. *J. Biol. Chem*. 1974; 249: 7130–39.
- [21] Habtermariam S, Gray AI, Waterman RG. A new anti-bacterial sesquiterpene from *Premna oligotricha*. *J. Nat. Prod*. 1993; 5: 140–43.

- [22] Heleno SA, Barros L, Martins A, Queiroz MJRP, Santos-Buelga C, Ferreira ICFR. Fruiting body, spores and in vitro produced mycelium of *Ganoderma lucidum* from Northeast Portugal: A comparative study of the antioxidant potential of phenolic and polysaccharidic extracts. *Food Res. Int.* 2012; 46: 135–40.
- [23] Hua X, Lei M, Ding J, Han Q, Hu G, Xiao M. Pathological and biochemical alterations of astrocytes in ovariectomized rats injected with D-galactose: a potential contribution to Alzheimer's disease processes. *Exp Neurol.* 2008; 210: 709-18
- [24] IACUC. International Animal Care and Use Committee Lab Animal. 2010; 39(6)
- [25] Ilahi I, Samar S, Khan I, Ahmad I. In vitro antioxidant activities of four medicinal plants on the basis of DPPH free radical scavenging. *Pak. J. Pharm. Sci.* 2013; 26 (5): 949-52
- [26] JollowDJ, Michell JR, Zampaglionic, Gillete JR. Bromoibenzene-induced Liver necrosis: Protective role of glutathione and evidence for 3,4- Bromobenzene oxide as hepatotoxic metabolite. *Pharmacology.*1974;11:151-69
- [27] Kalauni SK, Choudhary MI, Khalid A, Manandhar MD, Shaheen F, Atta-ur-Rahman, Gewali M. New cholinesterase inhibiting steroidal alkaloids from the leaves of *Sarcococca coriacea* of Nepalese origin. *Chem Pharm Bull.* 2002; 50: 1423-26
- [28] Krishnamurthy P, Wadhvani A. Antioxidant Enzymes and Human Health. InTech open access chapter distributed under the terms of the Creative Commons Attribution License. 2012; 13-15
- [29] Lahiri DK, Farlow MR, Greig NH, Sambamurti K. Current drug targets for Alzheimer's disease treatment. *Drug Dev. Res.* 2002; 56: 267–21.
- [30] Liu IX, Durham DG, Richards ME. Vancomycin resistance reversal in *Enterococci* by flavonoids. *J. Pharm. Pharmacol.* 2001; 53: 129-32.
- [31] Mahdy K, Shaker O, Wafay H, Nassar Y, Hassan H, Hussein A. Effect of some medicinal plant extracts on the oxidative stress status in Alzheimer's disease induced in rats. *Eur Rev Med Pharmacol Sci.* 2012; 16(3): 31-42.
- [32] Maysaa AH, Eman HH, Naser JK, Dhefaf ZA, Ali HA, Haider KZ. Ameliorative Effect of *Curcuma longa* L. Rhizomes against Biochemical Toxicity Induced by Dichlorvos in Female Albino Rats. *Journal of Chemical and Pharmaceutical Sciences* 2016; 9(3):1098-1106
- [33] Merghem M, Dahamna S, Khennouf S. In-vivo Antioxidant Activity of *Rutamontana* L. Extracts. *J. Mater. Environ. Sci.*, 2019; 10(5): 470-77
- [34] Misra HP, FridovichI. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem.* 1972; 247:3170-5.
- [35] Mohandas J, Marshal JJ, Duggin GG, Horvath JS and Tiller DG. Low activities of glutathione-related enzymes as factors in the genesis of urinary bladder cancer. *Cancer. Res.* 1984; 44(11): 5086-91
- [36] Murraya AP, Faraonia MB, Castroa MJ, Alzaa NP, Cavallaro V. Natural AChE Inhibitors from Plants and their Contribution to Alzheimer's Disease Therapy. *Cur. Neuropharm.*, 2013; 11: 388-413
- [37] Naazeri S, Rostamian M, Yaghmaei B, Hedayati M. Fluorimetry as a Simple and Sensitive Method for Determination of Catalase Activity. *Zahedan Journal of Research in Medical Sciences.* 2014; 16: 64-67.
- [38] Nandy S, Paul HS, Barman NR, Chakraborty. In vitro evaluation of antioxidant activity of *Leucas plukenetii* (Roth) Spreng. *Asian Journal of Plant Sci. & Research.* 2012; 2: 254-62.

- [42] Nilima SR, Hande SM. Estimation of Phytochemical Content and Antioxidant Activity of Some Selected Traditional Indian Medicinal Plants. *Indian J Pharm Sci.* 2011; 73 (2): 146-51
- [43] Nwidu LL, Elmorsy E, Thornton J, Wijamunige B, Wijesekara A, Tarbox R. Anti-acetylcholinesterase activity and antioxidant properties of extracts and fractions of *Carpolobia lutea*. *Pharmaceutical Biology.* 2017; 55(1): 1875-83
- [44] Ojo OA, Oyinloye BE, Ajiboye BO, Ojo AB, Musa H, Olarewaju OI. Dichlorvos-induced oxidative stress in rat brain: Protective effects of the ethanolic extract of *Alstoniaboonei* stem bark. *Asian J Pharm.* 2014a; 8:216-21.
- [45] Ojo OA, Oyinloye BE, Ajiboye BO, Ojo AB, Akintayo CO and Okezie B: Dichlorvos Induced Nephrotoxicity in Rat Kidney: Protective Effects of *AlstoniaBoonei* Stem Bark Extract. *Int J Pharmacognosy* 2014b; 1(7) 429-437
- [46] Oseni OA, Williams OD. In-vitro compositional investigations of antioxidants, phytochemicals, nutritional and minerals in the fruit of *Kigeliaafricana* (Lam.) Benth. *International J. of Contemporary Research and Review.* 2018; 9(8) 20259-68
- [47] Ruch RJ, Cheng SJ, Klaunig JE. *Carcinogenesis, Rely. Chim. Acta.* 1989; 10, 1003-08
- [48] Sato MH, Tsuchiya T, Miyazaki M, Ohyama T, Linuma M. Antibacterial activity of flavanostilbens against methicillin-resistant *Staphylococcus aureus*. *Lett. Applied Microbiol.* 1995; 21: 219-22.
- [49] Shirwaikar A, Shirwaikar A, Rajendran K, Punitha ISJ. In Vitro Antioxidant Studies on the Benzyl Tetra Isoquinoline Alkaloid Berberine. *Biol. Pharm. Bull.* 2006; 29(9): 1906-10
- [50] Sofowora A: *Medicinal Plants and Traditional Medicinal in Africa.* 2nd Ed. Sunshine House, Ibadan, Nigeria: Spectrum Books Ltd; *Screening Plants for Bioactive Agents;* 1993; 134-56.
- [51] Sigma-Aldrich 2013. 'Technical Bulletin' Sigma-Aldrich Co. LLC, USA.
- [52] www.sigmaldrich.com
- [53] Singleton VL, Rossi JA. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am J Enol Vitic.* January 1965;16: 144-58.
- [54] Sung SY, Kang SY, Lee KY, Park MJ, Kim JH, Park JH, *et al.* (+)-a-Viniferin, a stilbene trimer from *Carangachamlague* inhibits acetylcholinesterase. *Biol Pharm Bull.* 2002; 25: 125-27
- [55] TMI. *The Merck Index. An Encyclopedia of Chemicals, Drugs, and Biologicals.* 11th ed. Ed. S. Budavari. Merck and Co. Inc., Rahway, NJ. 1989.
- [56] Trease GE, Evans WC: *Pharmacognosy.* 15th Ed. London: Saunders Publishers; 2002; 342-93.
- [57] Utley HG, Bernheim F, Hochstein P. Effect of sulfhydryl reagents on peroxidation in
- [58] microsomes. *Arch. Biochem. Biophys.* 1967; 118: 29
- [59] Vaghasiya Y, Dave R, Chanda S. Phytochemical Analysis of Some Medicinal Plants from Western Region of India. *Research Journal of Medicinal Plants.* 2011; 5(5): 567-76.
- [60] Volpicelli-Daley LA, Duysen EG, Lockridge O, Levey A. Altered hippocampal muscarinic receptors in acetylcholinesterase-deficient mice. *Ann Neurol* 2003; 53: 788-96
- [61] Von Euler HV, Josephson K. *European Journal of Organic Chemistry - Eur J. Org Chem.*
- [62] 1972; 452(1): 158-81.

- [63] Winterbourne CC, Hawkins RE, Brain M Carrel RW. The estimation of red cell superoxide dismutase activity. *J. Lab.chem.Med.* 1975; 85: 337-41.
- [64] Yadav P, Jadhav SE, Kumar V, Kaul KK, Pant SC, Flora SJS. Protective efficacy of 2-PAMCl, atropine and curcumin against dichlorvos induced toxicity in rats. *Interdiscip. Toxicol.* 2012; 5(1): 1-8
- [65] Zhishen J, Mengcheng T, Jianming W. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry.* 1999; 64: 555-59